

The Hilbert scheme of the diagonal in a product of projective spaces

Dustin Cartwright and Bernd Sturmfels

Abstract

The diagonal in a product of projective spaces is cut out by the ideal of 2×2 -minors of a matrix of unknowns. The multigraded Hilbert scheme which classifies its degenerations has a unique Borel-fixed ideal. This Hilbert scheme is generally reducible, and its main component is a compactification of $\mathrm{PGL}(d)^n / \mathrm{PGL}(d)$. For $n = 2$ we recover the manifold of complete collineations. For projective lines we obtain a novel space of trees that is irreducible but singular. All ideals in our Hilbert scheme are radical. We also explore connections to affine buildings and Deligne schemes.

1 Introduction

Multigraded Hilbert schemes parametrize families of ideals in a polynomial ring that share the same Hilbert function with respect to some grading by an abelian group [8]. We are interested in the following particular case. Let $X = (x_{ij})$ be a $d \times n$ -matrix of unknowns. We fix the polynomial ring $K[X]$ over a field K with the $\mathbb{Z}^n$ -grading by column degree, i.e. $\deg(x_{ij}) = e_j$. In this grading, the Hilbert function of the polynomial ring $K[X]$ equals

$$\mathbb{N}^n \rightarrow \mathbb{N}, (u_1, \dots, u_n) \mapsto \prod_{i=1}^n \binom{u_i + d - 1}{d - 1}.$$

We study the multigraded Hilbert scheme $H_{d,n}$, which parametrizes $\mathbb{Z}^n$ -homogeneous ideals I in $K[X]$ such that $K[X]/I$ has the Hilbert function

$$\mathbb{N}^n \rightarrow \mathbb{N}, (u_1, \dots, u_n) \mapsto \binom{u_1 + u_2 + \dots + u_n + d - 1}{d - 1}. \quad (1)$$

The key example is the ideal $I_2(X)$ that is generated by the 2×2 -minors of X , and whose quotient is indeed $\mathbb{Z}^n$ -graded with Hilbert function (1). The Hilbert scheme $H_{d,n}$ has the following geometric interpretation. Each $\mathbb{Z}^n$ -homogeneous ideal in $K[X]$ specifies a subscheme of the product of projective spaces $(\mathbb{P}^{d-1})^n = \mathbb{P}^{d-1} \times \dots \times \mathbb{P}^{d-1}$. The subscheme specified by the ideal $I_2(X)$ is the diagonal embedding of $\mathbb{P}^{d-1}$ in $(\mathbb{P}^{d-1})^n$. Our Hilbert scheme $H_{d,n}$ is a natural parameter space for degenerations of this diagonal in $(\mathbb{P}^{d-1})^n$.

The results obtained in this paper are as follows. In Section 2 we prove that all ideals I in $H_{d,n}$ are radical and Cohen-Macaulay. This result is derived by identifying a distinguished Borel-fixed ideal Z with these properties. It confirms a conjecture on multilinear Gröbner bases made by Conca in [5].

In Section 3 we show that $I_2(X)$ and one of its initial monomial ideals are smooth points on $H_{d,n}$. The irreducible component containing these points is an equivariant compactification of the homogeneous space G^n/G where $G = \mathrm{PGL}(d)$, and $G \subset G^n$ is the diagonal embedding. For $n = 2$ we recover Thaddeus' construction in [17] of the space of complete collineations. The relationship of our compactification of G^n/G to those constructed by Lafforgue in [12] will be discussed in Remark 3.8 and Example 4.10.

Section 4 is concerned with the case $d = 2$, and we regard its results to be the main contribution of this paper. We show that $H_{2,n}$ is irreducible, but singular for $n \geq 4$, and we determine its combinatorial structure. Each point in $H_{2,n}$ corresponds to a certain tree of projective lines. Among these are precisely $2^n(n+1)^{n-2}$ monomial ideals, one for each tree on $n+1$ unlabeled vertices and n labeled directed edges, and these form a graph.

In Section 5 we study the case $d = n = 3$. These are the smallest parameters for which the multigraded Hilbert scheme $H_{d,n}$ is reducible. We show that $H_{3,3}$ is the reduced union of seven irreducible components, with the main component of dimension 16 parametrizing degenerations of the diagonal in $\mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^2$. We list all monomial ideals on $H_{3,3}$ and their incidence relations.

Section 6 outlines a connection to convexity in affine buildings and tropical geometry. Extending previous results in [2, 11], we show how Gröbner degenerations on $H_{d,n}$ can be used to compute special fibers of Deligne schemes.

2 On a conjecture of Conca

Our plan is to derive Conca's Conjecture 4.2 in [5] from the following result.

Theorem 2.1. *All ideals I corresponding to points in $H_{d,n}$ are radical ideals.*

Proof. We may assume that K is an infinite field. Let $G = \mathrm{PGL}(d, K)$, the group of invertible $d \times d$ -matrices modulo scaling, let B be the Borel subgroup of images of upper triangular matrices in G , and let T be the algebraic torus of images of diagonal matrices in G . Then T^n is a maximal torus in G^n , and B^n is a Borel subgroup in G^n . We consider the action of these groups on the Hilbert scheme $H_{d,n}$. The T^n -fixed points of $H_{d,n}$ are the monomial ideals that have the same $\mathbb{Z}^n$ -graded Hilbert function as $I_2(X)$. It suffices to assume that I is such a monomial ideal because every other ideal $J \in H_{d,n}$ can be degenerated to a monomial ideal $I = \mathrm{in}(J)$ via Gröbner bases, and if $\mathrm{in}(J)$ is radical then so is J .

We can further assume that I is *Borel-fixed*, which means that I is fixed under the action of B^n . Indeed, if $A_1, A_2, \dots, A_n$ are generic matrices of G then we replace the ideal I first by its image $I' = (A_1, A_2, \dots, A_n) \circ I$, and then by the initial monomial ideal $\mathrm{in}(I')$. The ideal $\mathrm{in}(I') = \mathrm{gin}(I)$ is the *multigraded generic initial ideal*. The same approach as in [6, §15.9.2] shows that $\mathrm{gin}(I)$ is Borel-fixed. Moreover, if $\mathrm{gin}(I)$ is radical then so is I . Hence, it suffices to show that every Borel-fixed ideal I in $H_{d,n}$ is a radical ideal.

Our result will be a direct consequence of the following two claims:

Claim 1: *There is precisely one Borel-fixed ideal Z in $H_{d,n}$.*

Claim 2: *The unique Borel-fixed ideal Z is radical.*

We first describe the ideal Z and then prove that it has these properties. Let u be any vector in the set

$$U = \{(u_1, \dots, u_n) \in \mathbb{Z}^n : 0 \leq u_i \leq d-1 \text{ and } \sum_i u_i = (n-1)(d-1)\}. \quad (2)$$

We write Z_u for the ideal generated by all unknowns x_{ij} with $i \leq u_j$ and $1 \leq j \leq n$. This is a Borel-fixed prime monomial ideal. Consider the intersection of the prime ideals Z_u :

$$Z := \bigcap_{u \in U} Z_u.$$

The monomial ideal Z is a radical and Borel-fixed. Each of its $\binom{d+n-2}{d-1}$ associated prime ideals Z_u has the same codimension $(n-1)(d-1)$.

We now apply Conca's results in [5, Section 5]. He showed that Z has the same Hilbert function as $I_2(X)$. Therefore, the ideal Z is the promised Borel-fixed ideal in $H_{d,n}$. More precisely, [5, Theorem 5.1] states that Z is precisely the generic initial ideal $\mathrm{gin}(I_2(X))$ of the ideal of 2×2 -minors.

We claim that Z is the only Borel-fixed monomial ideal in $H_{d,n}$. To show this, we apply results about the *multidegree* in [13, §8.5]. The multidegree of the prime ideal Z_u is the monomial $\mathbf{t}^u = t_1^{u_1} t_2^{u_2} \cdots t_n^{u_n}$. By [13, Theorem 8.44], Z_u is the only unmixed Borel-fixed monomial ideal having multidegree $\mathbf{t}^u$. By [13, Theorem 8.53], the Borel-fixed ideal $Z = \cap_u Z_u$ has the multidegree

$$\mathcal{C}(k[X]/Z; \mathbf{t}) = \sum_{u \in U} \mathbf{t}^u = \sum_{u \in U} t_1^{u_1} t_2^{u_2} \cdots t_n^{u_n}. \quad (3)$$

Since the multidegree of a homogeneous ideal is determined by its Hilbert series [13, Claim 8.54], we conclude that every ideal $I \in H_{d,n}$ has multidegree (3). Now, suppose that $I \in H_{d,n}$ is Borel-fixed. Since I is monomial, each minimal primary component contributes at most one term $\mathbf{t}^u$ to the multidegree (3). Thus, by [13, Theorem 8.53], the minimal primes of I are precisely the prime ideals Z_u where u runs over the elements of U . This implies $I \subseteq \sqrt{I} = Z$. However, since I and Z have the same Hilbert function in a positive grading, we conclude that $I = Z$, as desired. $\square$

Remark 2.2. Our proof of Theorem 2.1 was based on an idea that was suggested to us by Michel Brion. In [4], Brion proves that for any multiplicity-free subvariety of a flag variety, such as the diagonal in a product of projective spaces, there exists a flat degeneration to a reduced union of Schubert varieties, which is our Z . Although Theorem 2.1 only applies to the special case of the diagonal in a product of projective spaces, it establishes reducedness not just for *some* degeneration but for *any* ideal with the same multigraded Hilbert function. Our proof combined the nice argument from [4] with the explicit description of the Borel-fixed monomial ideal given by Conca in [5]. $\square$

We now come to the question asked by Conca in [5, Conjecture 1.1]. Given any $d \times d$ -matrices $A_1, A_2, \dots, A_n$ with entries in K , we apply them individually to the n columns of the matrix $X = (x_{ij})$, form the ideal of 2×2 -minors of the resulting $d \times n$ -matrix, and then take the initial monomial ideal

$$\text{in}((A_1, \dots, A_n) \circ I_2(X)) \quad (4)$$

with respect to some term order. Conca conjectures that (4) is always a squarefree monomial ideal. He proves this for generic A_i by showing that (4) equals the Borel-fixed ideal Z constructed above. Theorem 2.1 implies the same conclusion under the much weaker hypothesis that the A_i are invertible.

Corollary 2.3. *For any invertible $d \times d$ -matrices $A_1, \dots, A_n$ and any term order on $K[X]$, the monomial ideal $\text{in}((A_1, \dots, A_n) \circ I_2(X))$ is squarefree.*

Proof. Applying invertible matrices A_i to $I_2(X)$ corresponds to taking the orbit of $I_2(X)$ under the action of G^n on $H_{d,n}$. Therefore, (4) is a monomial ideal that lies in $H_{d,n}$. By Theorem 2.1, it is radical and hence squarefree. $\square$

Corollary 2.3 implies [5, Conjecture 4.2]. At present, we do not know how to prove Conca's stronger conjecture to the effect that Corollary 2.3 holds without the hypothesis that the matrices A_i are invertible [5, Conjecture 1.1]. One idea is to extend our study to multigraded Hilbert schemes on $K[X]$ whose defining Hilbert function is bounded above by (1).

Corollary 2.4. *The multigraded Hilbert scheme $H_{d,n}$ is connected.*

Proof. All ideals in $H_{d,n}$ can be connected to their common generic initial ideal Z by Gröbner degenerations. $\square$

In what follows we take a closer look at the combinatorics of the ideal Z .

Proposition 2.5. *The ideal Z is generated by all monomials $x_{i_1 j_1} x_{i_2 j_2} \dots x_{i_k j_k}$ where $1 \leq k-1 \leq i_1, i_2, \dots, i_k \leq d-1$, $j_1 < j_2 < \dots < j_k$, and $i_1 + i_2 + \dots + i_k \leq d(k-1)$. The maximum degree of a minimal generator is $\min(d, n)$.*

Proof. This is the description of the ideal Z given by Conca [5, §5]. $\square$

All ideals I in $H_{d,n}$ share the same Hilbert series in the ordinary $\mathbb{Z}$ -grading,

$$\sum_{r=0}^{\infty} \dim_K(K[X]/I)_r \cdot z^r = \frac{h(z)}{(1-z)^{n+d-1}}.$$

The h -polynomial in the numerator can be seen from the ideal of 2×2 -minors:

$$h(z) = \sum_{i=0}^{\min(d-1, n-1)} \binom{d-1}{i} \cdot \binom{n-1}{i} \cdot z^i.$$

Note that $h(1) = \binom{n+d-2}{d-1}$ is the common scalar degree of the ideals in $H_{d,n}$.

Corollary 2.6. *Every ideal I in $H_{d,n}$ is Cohen-Macaulay.*

Proof. Since Z is the common generic initial ideal of all ideals I , it suffices to show that the Borel-fixed ideal Z is Cohen-Macaulay. Let Δ_Z denote the $(n + d - 2)$ -dimensional simplicial complex corresponding to Z . The vertices of Δ_Z are the dn matrix entries x_{ij} , and its facets are the $\binom{n+d-2}{d-1}$ sets $F_u = \{x_{ij} : i > u_j\}$ which are complementary to the prime ideals Z_u . We order the facets F_u according to the lexicographic order on the vectors u .

We claim that this ordering of the facets is a *shelling* of Δ_Z . Since the Stanley-Reisner ring of a shellable simplicial complex is Cohen-Macaulay [16, Theorem III.2.5], this will imply Corollary 2.6. To verify the shelling property we must show that every facet F_u has a unique subset η_u such that the faces of F_u not containing η_u are exactly those appearing as a face of an earlier facet. If this condition holds then the h -polynomial can be read off from the shelling as follows:

$$h(z) = \sum_{u \in U} z^{\#\eta_u}.$$

The unique subset of the facet F_u with these desired properties equals

$$\eta_u = \{x_{ij} : j > 1 \text{ and } i = u_j + 1 < d\}.$$

Indeed, suppose G is a face common to F_u and $F_{u'}$ for some $u' < u$. Then $u'_j > u_j$ for some $j > 1$, so G does not contain $x_{u_j+1,j} \in \eta_u$. Conversely, suppose that G is a face of F_u which does not contain η_u , and let $x_{u_j+1,j}$ be any element of $\eta_u \setminus G$. Since $j > 1$,

$$F_{u+e_j-e_1} = F_u \setminus \{x_{u_j+1,j}\} \cup \{x_{u_1,1}\}$$

is a facet of Δ_Z which contains G and which comes earlier in our ordering. $\square$

Remark 2.7. The shellability of Δ_Z was mentioned in [5, Remark 5.12] but no details were given there. It would be interesting to know whether the simplicial complex Δ_I of every monomial ideal I in $H_{d,n}$ is shellable.

3 Group completions

In this section we relate our multigraded Hilbert scheme to classical constructions in algebraic geometry. For $n = 2$ we recover the space of complete collineations and its GIT construction due to Thaddeus in [17]. Brion [3] extended Thaddeus' work to the diagonal $X \hookrightarrow X \times X$ of any rational projective homogeneous variety X . While the present study is restricted to the

case $X = \mathbb{P}^{d-1}$, we believe that many of our results will extend to $X \hookrightarrow X^n$ in Brion's setting.

Proposition 3.1. *There is an injective morphism from the multigraded Hilbert scheme $H_{d,n}$ to a connected component of the Grothendieck Hilbert scheme of subschemes of $(\mathbb{P}^{d-1})^n$.*

Proof. By [8, §4], there is a natural morphism from the multigraded Hilbert scheme $H_{d,n}$ to the Grothendieck Hilbert scheme. Theorem 2.1 shows that every point in $H_{d,n}$ corresponds to a radical ideal $I \subset K[X]$. Furthermore, the Hilbert function tells us that the ideal $I + \langle x_{1j}, \dots, x_{dj} \rangle$ has (affine) dimension $d + n - 2$, but by Corollary 2.6, I is pure of dimension $d + n - 1$, so no associated prime of I contains $\langle x_{1j}, \dots, x_{dj} \rangle$. Thus, I is uniquely determined by the subscheme it defines in $(\mathbb{P}^{d-1})^n$, so the morphism is injective. $\square$

Remark 3.2. We do not know whether the morphism in Proposition 3.1 is an immersion, nor whether it is always surjective.

Recall that the G^n -action on $H_{d,n}$ transforms ideals as follows:

$$I \mapsto (A_1, A_2, \dots, A_n) \circ I. \quad (5)$$

If $A_1 = A_2 = \dots = A_n$ then the ideal I is left invariant, so the stabilizer of any point $I \in H_{d,n}$ contains the diagonal subgroup $G = \{(A, A, \dots, A)\}$ of G^n . Moreover, the stabilizer of the determinantal ideal $I_2(X)$ is precisely the diagonal subgroup G . We write $\overline{G^n/G}$ for the closure of $G^n \circ I_2(X)$ in the Hilbert scheme $H_{d,n}$.

Theorem 3.3. *The subscheme $\overline{G^n/G}$ is an irreducible component of $H_{d,n}$. It is a compactification of the homogeneous space G^n/G , so it has dimension $(d^2 - 1)(n - 1)$.*

This theorem can be deduced from Proposition 3.1 using standard algebraic geometry arguments concerning the tangent sheaf of Grothendieck's Hilbert scheme. What we present below is a more detailed combinatorial proof based on the identification of an explicit smooth point in Lemma 3.4.

Proof. Clearly, the dimension of the tangent space at $I_2(X)$ is at least $(d^2 - 1)(n - 1)$, the dimension of G^n/G . By semi-continuity, it is bounded above by the tangent space dimension of $H_{d,n}$ at any initial monomial ideal in $(I_2(X))$. In Lemma 3.4 below, we identify a particular initial ideal for which this

dimension equals $(d^2 - 1)(n - 1)$. From this we conclude that the tangent space of $H_{d,n}$ at $I_2(X)$ has dimension $(d^2 - 1)(n - 1)$. This is precisely the dimension of the orbit closure $\overline{G^n/G}$ of $I_2(X)$. We also conclude that $I_2(X)$ is a smooth point of $H_{d,n}$, and that the unique irreducible component of $H_{d,n}$ containing $I_2(X)$ is the compactified space $\overline{G^n/G}$. $\square$

Let M denote the ideal generated by the quadratic monomials $x_{ik}x_{jl}$ for all $1 \leq i < j \leq d$ and $1 \leq k < l \leq n$. We call M the *chain ideal*, because its irreducible components correspond to chains in the grid from x_{1n} to x_{d1} . It is a point in $\overline{G^n/G} \subset H_{d,n}$ since $M = \text{in}(I_2(X))$ in the lexicographic order.

Lemma 3.4. *The tangent space of the multigraded Hilbert scheme $H_{d,n}$ at the chain ideal M has dimension $(d^2 - 1)(n - 1)$.*

Proof. We claim that the following three classes ρ , σ and τ form a basis for the tangent space $\text{Hom}_{K[X]}(M, K[X]/M)_0$:

Class ρ : For each triple of indices (i, j, l) with $1 \leq i < j \leq d$ and $1 < l \leq n$ we define a $K[X]$ -module homomorphism $\rho_{ijl}: M \rightarrow K[X]/M$ by setting

$$\begin{aligned} \rho_{ijl}(x_{hk}x_{jl}) &= x_{hk}x_{il} \quad \text{whenever } i \leq h < j \text{ and } k < l, \text{ and} \\ \rho_{ijl}(m) &= 0 \quad \text{for all other minimal generators } m \text{ of } M. \end{aligned}$$

Class σ : For each triple of indices (i, j, k) with $1 \leq i < j \leq d$ and $1 \leq k < n$ we define a $K[X]$ -module homomorphism $\sigma_{ijk}: M \rightarrow K[X]/M$ by setting

$$\begin{aligned} \sigma_{ijk}(x_{ik}x_{hl}) &= x_{jk}x_{hl} \quad \text{whenever } i < h \leq j \text{ and } k < l, \text{ and} \\ \sigma_{ijk}(m) &= 0 \quad \text{for all other minimal generators } m \text{ of } M. \end{aligned}$$

Class τ : For each pair of indices (i, k) with $1 \leq i < d$ and $1 \leq k < n$ we define a $K[X]$ -module homomorphism $\tau_{ik}: M \rightarrow K[X]/M$ by setting

$$\begin{aligned} \tau_{ik}(x_{i,k}x_{i+1,k+1}) &= x_{i,k+1}x_{i+1,k} \quad \text{and,} \\ \tau_{ik}(m) &= 0 \quad \text{for all other minimal generators } m \text{ of } M. \end{aligned}$$

The above $K[X]$ -linear maps are $\mathbb{Z}^d$ -homogeneous of degree zero, and they are clearly linearly independent over K . There are $(n - 1)(d - 1)$ maps in the class τ , and there are $(n - 1)\binom{d}{2}$ each in the classes ρ and σ . This adds up to the required total number of $(d^2 - 1)(n - 1) = (n - 1)(d - 1) + 2(n - 1)\binom{d}{2}$.

It remains to be seen that every $\mathbb{Z}^d$ -graded $K[X]$ -module homomorphism from M to $K[X]/M$ of degree zero is a K -linear combination of the above. Suppose that $\phi: M \rightarrow K[X]/M$ is a module homomorphism. Then, for $i < j$ and $k < l$, we can uniquely write $\phi(x_{ik}x_{jl})$ as a linear combination of monomials not in M . Furthermore, by subtracting appropriate multiples of ρ_{ijl} and σ_{ijk} , we can assume that the monomials in the linear combination do not include $x_{ik}x_{il}$ or $x_{jk}x_{jl}$. Suppose that for some $n \leq m$, the coefficient of $x_{mk}x_{nl}$ is some non-zero $\alpha \in K$. Either $i < m$ or $n < j$, and the two cases are symmetric under reversing the order of both the column indices and the row indices, so we assume the former. For any o such that $n \leq o \leq m$ and $i < o$, the syzygies imply

$$\alpha x_{o,k+1}x_{mk}x_{nl} + \cdots = x_{o,k+1}\phi(x_{ik}x_{jl}) = x_{jl}\phi(x_{ik}x_{o,k+1}).$$

Since the first term is non-zero in $K[X]/M$, the monomial must be divisible by x_{jl} . Thus, either $j = n$, or both $j = o$ and $l = k + 1$. In the first case, taking $o = m$, and using the assumption that the coefficient of $x_{mk}x_{m,k+1}$ in $\phi(x_{ik}x_{o,k+1})$ is zero, we get a contradiction. In the second case, if $j \neq n$, then we must only have one choice of o and this forces $i = n = m - 1$. Therefore, ϕ is a linear combination of homomorphisms of class τ , and thus the classes of ρ , σ , and τ span the tangent space at M . $\square$

Corollary 3.5. *The chain ideal M is a smooth point on $H_{d,n}$. The unique irreducible component of $H_{d,n}$ containing M is the completion $\overline{G^n}/G$.*

We now turn to the case $n = 2$ which is well-studied in the literature. The compactification $\overline{G^2}/G$ is the classical *space of complete collineations*, which was investigated by Thaddeus in [17]. In fact, we have:

Corollary 3.6. *The multigraded Hilbert scheme $H_{d,2}$ is smooth and irreducible. It coincides with the space of complete collineations: $H_{d,2} = \overline{G^2}/G$.*

Proof. Up to relabeling, the chain ideal is the only monomial ideal in $H_{d,2}$. This point is smooth by Lemma 3.5, and hence $H_{d,2}$ is smooth. Since it is connected by Corollary 2.4, we conclude that $H_{d,2}$ is also irreducible. The results in [17] show that the Grothendieck Hilbert scheme is isomorphic to the space of complete collineations, and in particular smooth and irreducible. Thus, the morphism in Proposition 3.1 is an isomorphism between $H_{d,2}$ and the space of complete collineations. $\square$

The representation of $\overline{G^2/G}$ as a multigraded Hilbert scheme $H_{d,2}$ gives rise to nice polynomial equations for the space of complete collineations. Namely, each ideal I in $H_{d,2}$ is generated by $\binom{d}{2}$ equations of degree $(1, 1)$. As there are d^2 monomials in $K[X]_{(1,1)}$, this describes an embedding of $H_{d,2}$ into the Grassmannian $\text{Gr}(\binom{d}{2}, d^2)$. The subscheme $H_{d,2}$ of this Grassmannian is cut out by the determinantal equations which are derived by requiring that the ideal I has the correct number of first syzygies in degrees $(1, 2)$ and $(2, 1)$.

Example 3.7 (Equations defining $H_{3,2}$). We shall realize the 8-dimensional manifold $H_{3,2}$ as a closed subscheme of the 18-dimensional Grassmannian $\text{Gr}(3, 9)$, by giving explicit equations in the 84 Plücker coordinates. Our equations furnish an explicit projective embedding for Thaddeus' GIT construction [17] which is reviewed further below. Fix a 3×9 -matrix of unknowns

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{21} & a_{22} & a_{23} & a_{31} & a_{32} & a_{33} \\ b_{11} & b_{12} & b_{13} & b_{21} & b_{22} & b_{23} & b_{31} & b_{32} & b_{33} \\ c_{11} & c_{12} & c_{13} & c_{21} & c_{22} & c_{23} & c_{31} & c_{32} & c_{33} \end{bmatrix}.$$

Consider the ideal I generated by the three bilinear polynomials in the vector

$$A \cdot (x_{11}x_{12}, x_{11}x_{22}, x_{11}x_{32}, x_{21}x_{12}, x_{21}x_{22}, x_{21}x_{32}, x_{31}x_{12}, x_{31}x_{22}, x_{31}x_{32})^T.$$

The condition for I to be a point in $H_{3,2}$ is equivalent to the condition that the rows of the following two 9×18 -matrices are linearly dependent:

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & a_{21} & a_{22} & a_{23} & a_{31} & a_{32} & a_{33} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ b_{11} & b_{12} & b_{13} & b_{21} & b_{22} & b_{23} & b_{31} & b_{32} & b_{33} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ c_{11} & c_{12} & c_{13} & c_{21} & c_{22} & c_{23} & c_{31} & c_{32} & c_{33} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & a_{11} & a_{12} & a_{13} & 0 & 0 & 0 & a_{21} & a_{22} & a_{23} & a_{31} & a_{32} & a_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & b_{11} & b_{12} & b_{13} & 0 & 0 & 0 & b_{21} & b_{22} & b_{23} & b_{31} & b_{32} & b_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & c_{11} & c_{12} & c_{13} & 0 & 0 & 0 & c_{21} & c_{22} & c_{23} & c_{31} & c_{32} & c_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & a_{11} & a_{12} & a_{13} & 0 & 0 & 0 & a_{21} & a_{22} & a_{23} & a_{31} & a_{32} & a_{33} \\ 0 & 0 & 0 & 0 & 0 & 0 & b_{11} & b_{12} & b_{13} & 0 & 0 & 0 & b_{21} & b_{22} & b_{23} & b_{31} & b_{32} & b_{33} \\ 0 & 0 & 0 & 0 & 0 & 0 & c_{11} & c_{12} & c_{13} & 0 & 0 & 0 & c_{21} & c_{22} & c_{23} & c_{31} & c_{32} & c_{33} \end{bmatrix}$$

$$\begin{bmatrix} a_{11} & a_{21} & a_{31} & a_{12} & a_{22} & a_{32} & a_{13} & a_{23} & a_{33} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ b_{11} & b_{21} & b_{31} & b_{12} & b_{22} & b_{32} & b_{13} & b_{23} & b_{33} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ c_{11} & c_{21} & c_{31} & c_{12} & c_{22} & c_{32} & c_{13} & c_{23} & c_{33} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & a_{11} & a_{21} & a_{31} & 0 & 0 & 0 & a_{12} & a_{22} & a_{32} & a_{13} & a_{23} & a_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & b_{11} & b_{21} & b_{31} & 0 & 0 & 0 & b_{12} & b_{22} & b_{32} & b_{13} & b_{23} & b_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & c_{11} & c_{21} & c_{31} & 0 & 0 & 0 & c_{12} & c_{22} & c_{32} & c_{13} & c_{23} & c_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & a_{11} & a_{21} & a_{31} & 0 & 0 & 0 & a_{12} & a_{22} & a_{32} & a_{13} & a_{23} & a_{33} \\ 0 & 0 & 0 & 0 & 0 & 0 & b_{11} & b_{21} & b_{31} & 0 & 0 & 0 & b_{12} & b_{22} & b_{32} & b_{13} & b_{23} & b_{33} \\ 0 & 0 & 0 & 0 & 0 & 0 & c_{11} & c_{21} & c_{31} & 0 & 0 & 0 & c_{12} & c_{22} & c_{32} & c_{13} & c_{23} & c_{33} \end{bmatrix}$$

These two matrices are obtained by multiplying the generators of I with the entries in the two columns of $X = (x_{ij})$ respectively. This results in nine polynomials of bidegree $(2, 1)$ and nine polynomials of bidegree $(1, 2)$, each having 18 terms. These two sets of polynomials must be linearly dependent because each $I \in H_{3,2}$ has its first syzygies in these two bidegrees.

The 84 maximal minors of the matrix A are the Plücker coordinates $p_{i_1 i_2, j_1 j_2, k_1 k_2}$ on the Grassmannian $\text{Gr}(3, 9)$, where the indices run from 1 to 3. Using Laplace expansion, we write each 9×9 -minor of the two matrices as a cubic polynomial in these Plücker coordinates. The condition that the matrices have rank at most eight translates into a system of homogeneous cubic polynomials in the 84 unknowns $p_{i_1 i_2, j_1 j_2, k_1 k_2}$, and these cubics define the space of complete collineations, $\overline{G^2/G} = H_{3,2}$, as a subscheme of $\text{Gr}(3, 9)$.

Thaddeus [17] realizes $H_{3,2}$ as the (Chow or GIT) quotient of the Grassmannian $\text{Gr}(3, 6)$ by the one-dimensional subtorus of $(K^*)^6$ given by the diagonal matrices with entries $(t, t, t, t^{-1}, t^{-1}, t^{-1})$. We can see this in our equations as follows. Let $U = (u_{ij})$ and $V = (v_{ij})$ be 3×3 -matrices of unknowns. Each point in $\text{Gr}(3, 6)$ is represented as the row space of the 3×6 -matrix $[U, V]$. The group $G^2 = \text{PGL}(3) \times \text{PGL}(3)$ acts on $H_{3,2}$ by translating the distinguished point $I_2(X)$ to the ideal generated by the three quadrics

$$\begin{aligned} & (u_{i1}x_{11} + u_{i2}x_{21} + u_{i3}x_{31})(v_{j1}x_{12} + v_{j2}x_{22} + v_{j3}x_{32}) \\ - & (u_{j1}x_{11} + u_{j2}x_{21} + u_{j3}x_{31})(v_{i1}x_{12} + v_{i2}x_{22} + v_{i3}x_{32}) \end{aligned} \quad \text{for } 1 \leq i < j \leq 3.$$

The entries of the corresponding 3×9 matrix A are

$$a_{i_1 i_2} = u_{1i_1}v_{2i_2} - u_{2i_1}v_{1i_2}, \quad b_{j_1 j_2} = u_{1j_1}v_{3i_2} - u_{3j_1}v_{1j_2}, \quad c_{k_1 k_2} = u_{2k_1}v_{3i_2} - u_{3k_1}v_{2k_2}.$$

Writing u_μ for the μ -th column of the matrix U and v_ν for the ν -th column of V , this translates into the following parametric representation of $H_{3,2}$:

$$p_{i_1 i_2, j_1 j_2, k_1 k_2} = \det[u_{i_1}, v_{i_2}, u_{j_1}] \det[v_{j_2}, u_{k_1}, v_{k_2}] - \det[u_{i_1}, v_{i_2}, v_{j_2}] \det[u_{j_1}, u_{k_1}, v_{k_2}].$$

These are quadratic polynomials in the Plücker coordinates on $\text{Gr}(3, 6)$. They are invariant under Thaddeus' torus action and antisymmetric under swapping each of the three index pairs. Note that of the 84 polynomials only 12 are actually Plücker binomials. Of the others, 6 are zero (for example, $p_{11,21,31} = 0$) and 66 are Plücker monomials (for example, $p_{11,21,32} = \det[u_1, v_1, u_2] \det[v_1, u_3, v_2]$). This resulting map $\text{Gr}(3, 6) \rightarrow \text{Gr}(3, 9)$ gives an embedding of Thaddeus' quotient $H_{3,2} = \text{Gr}(3, 6)/K^*$. The cubic relations on $\text{Gr}(3, 9)$ described above characterize the image of this embedding. $\square$

Remark 3.8. In the introduction of [12], Lafforgue describes the following compactification of G^n/G . We consider the $d \times d$ -minors of the $d \times (dn)$ -matrix $(A_1, A_2, \dots, A_n)$. For each minor there is a corresponding vector $\mathbf{i}$ in the set $D = \{(i_1, i_2, \dots, i_n) \in \mathbb{N}^n : i_1 + \dots + i_n = d\}$, namely, i_j is the number of columns of A_j occurring in that minor. We introduce a new unknown $t_{\mathbf{i}}$ for each $\mathbf{i} \in D$, and we multiply each minor by the corresponding unknown $t_{\mathbf{i}}$. The scaled minors parametrize a subvariety in an affine space of dimension

$$\binom{nd}{d} = \sum_{\mathbf{i} \in D} \binom{d}{i_1} \binom{d}{i_2} \cdots \binom{d}{i_n}.$$

This affine variety yields a projective variety $X_{d,n}$ which compactifies G^n/G :

$$G^n/G \hookrightarrow X_{d,n} \subset \prod_{\mathbf{i} \in D} \mathbb{P}^{\binom{d}{i_1} \binom{d}{i_2} \cdots \binom{d}{i_n} - 1}. \quad (6)$$

In light of [8, §2], we can identify $X_{d,n}$ with the partial multigraded Hilbert scheme $(H_{d,n})_D$ obtained by restricting $H_{d,n}$ to the subset of degrees $D \subset \mathbb{Z}^n$. Hence there is a natural morphism $H_{d,n} \rightarrow X_{d,n}$. This is an isomorphism for $d = 2$ and $n = 2$ but we do not know whether this is always the case. In general, $X_{d,n}$ is singular, and the main result of [12] is a combinatorial construction that replaces $X_{d,n}$ with another – less singular – model $\Omega_{d,n}$. Yet, as discussed in the erratum to [12], $\Omega_{d,n}$ is not smooth for $d, n \geq 4$. $\square$

4 Yet another space of trees

This section concerns the case $d = 2$. The Hilbert scheme $H_{2,n}$ parametrizes degenerations of the projective line in its diagonal embedding $\mathbb{P}^1 \hookrightarrow (\mathbb{P}^1)^n$. Our goal is to prove the following two theorems about the structure of $H_{2,n}$.

Theorem 4.1. *The multigraded Hilbert scheme $H_{2,n}$ is irreducible, so it equals the compactification $\overline{\mathrm{PGL}(2)^n/\mathrm{PGL}(2)}$. However, $H_{2,n}$ is singular for $n \geq 4$.*

Our second theorem explains why we refer to $H_{2,n}$ as a *space of trees*. The qualifier “yet another” has been prepended to emphasize that this is not the *space of phylogenetic trees*. The latter is familiar to algebraic geometers as a discrete model for $\overline{\mathcal{M}}_{0,n}$; see [10, Theorem 1.2] for a precise statement.

Following [1], there is a natural graph structure on any multigraded Hilbert scheme, including $H_{2,n}$. The vertices are the monomial ideals, and for every ideal in $H_{2,n}$ with precisely two initial monomial ideals there is an edge between the corresponding vertices. By [1, Theorem 11], this is precisely the induced subgraph on $H_{2,n}$ of the graph of all monomial ideals. We note that our graph is not a *GKM graph* in the sense of [7] because the T^n -fixed subvarieties corresponding to edges usually have dimension greater than one.

Theorem 4.2. *There are $2^n(n+1)^{n-2}$ monomial ideals in $H_{2,n}$, one for each tree on $n+1$ unlabeled vertices with n labeled directed edges. Two trees are connected by an edge on $H_{2,n}$ if they differ by one of the following operations:*

1. *Move any subset of the trees attached at a vertex to an adjacent vertex.*
2. *Swap two edges that meet at a bivalent vertex (preserving orientation).*

In this section we use the following notation for our matrix of variables:

$$X = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \\ y_1 & y_2 & \cdots & y_n \end{bmatrix}.$$

Thus $(x_i : y_i)$ are homogeneous coordinates on the i -th factor in our ambient space $(\mathbb{P}^1)^n$. The common Hilbert function (1) of all ideals I in $H_{2,n}$ equals

$$\mathbb{N}^n \rightarrow \mathbb{N}, (u_1, u_2, \dots, u_n) \mapsto u_1 + u_2 + \cdots + u_n + 1. \quad (7)$$

The unique Borel-fixed ideal in $H_{2,n}$ equals $Z = \langle x_i x_j : 1 \leq i < j \leq n \rangle$. Our first goal is to prove that $H_{2,n}$ is irreducible. This requires a combinatorial description of the subvarieties $V(I)$ of $(\mathbb{P}^1)^n$ corresponding to ideals $I \in H_{2,n}$. Note that each such subvariety is a reduced curve of multidegree $(1, 1, \dots, 1)$ in $(\mathbb{P}^1)^n$.

Lemma 4.3. *The variety $V(I) \subset (\mathbb{P}^1)^n$ defined by any ideal $I \in H_{2,n}$ is the reduced union of several copies of the projective line $\mathbb{P}^1$. For each factor of $(\mathbb{P}^1)^n$ there is exactly one component of $V(I)$ which is not constant along this factor, and for this component, the projection induces an isomorphism.*

Proof. Consider the projection from $V(I)$ onto the i -th factor of $(\mathbb{P}^1)^n$. We infer from the Hilbert function (7) that this projection is an isomorphism over an open subset of $\mathbb{P}^1$. Hence there exists a rational map from $\mathbb{P}^1$ to a unique component Y of $V(I)$. The fact that the projection $Y \rightarrow \mathbb{P}^1$ is a regular morphism implies that the curve Y is smooth. We conclude that the map $Y \rightarrow \mathbb{P}^1$ is an isomorphism. $\square$

Each component of $V(I)$ can be labeled by the factors onto which it maps isomorphically. We draw $V(I)$ as a set of intersecting lines, labeled with subsets of the factors. By Lemma 4.3, the labels form a partition of $\{1, 2, \dots, n\}$. Moreover, since $K[X]/I$ is Cohen-Macaulay, $V(I)$ is connected, and because only one component is non-constant along any factor, there is no cycle among its components. Hence our picture is an edge-labeled tree. We have the following converse to this description of the points of $H_{2,n}$.

Proposition 4.4. *Suppose that $Y \subset (\mathbb{P}^1)^n$ is a union of projective lines, which is connected and such that each factor of $(\mathbb{P}^1)^n$ has a unique projective line projecting isomorphically onto it. Then the radical ideal I defining Y is a point in $H_{2,n}$.*

Proof. We compute the Hilbert function and show that it coincides with (7). We proceed by induction on the number of components. If Y is irreducible then Y is the translate of the diagonal $\mathbb{P}^1$ in $(\mathbb{P}^1)^n$ with some $A_i \in \text{PGL}(2)$ acting on the i -th factor, and therefore $I = (A_1, \dots, A_n) \circ I_2(X)$ lies in $H_{2,n}$.

Now suppose Y is reducible, let Y_j be one of its components, and $F_j \subset \{1, \dots, n\}$ the index set of factors of $(\mathbb{P}^1)^n$ onto which Y_j maps isomorphically. The prime ideal I_j of Y_j is generated by linear forms of multidegrees $\{e_i : i \notin F_j\}$, and by the 2×2 -minors of a $2 \times |F_j|$ -matrix X_j which consists of the F_j columns of X acted on by some $A_i \in \text{PGL}(2)$. The Hilbert function of I_j is

$$u \mapsto 1 + \sum_{i \in F_j} u_i.$$

Since Y is a tree of projective lines, there exists a component Y_j which has only one point of intersection with the other components. Let $Y' \subset (\mathbb{P}^1)^n$

be the union of the other components and I' the radical ideal defining Y' . The ideal I_j of Y_j contains linear forms of degree e_i for every $i \notin F_j$, while I' contains linear forms of degree e_i for every $i \in F_j$. This implies that $I' + I_j$ is a homogenous prime ideal generated by linear forms. Its variety equals $Y' \cap Y_j$, and hence $I' + I_j$ has constant Hilbert function 1. We conclude

$$\begin{aligned} HF(I) &= HF(I' \cap I_j) = HF(I') + HF(I_j) - HF(I' + I_j) \\ &= \left(1 + \sum_{i \in F_j} u_i\right) + \left(1 + \sum_{i \notin F_j} u_i\right) - 1, \end{aligned}$$

which is the common Hilbert function (7) of all ideals in $H_{2,n}$. $\square$

Our discussion shows that each point in $H_{2,n}$ is characterized by the following data. First, there is a tree of projective lines $Y_j = \mathbb{P}^1$, labeled by the parts in a partition $\{1, \dots, n\} = \cup_j F_j$. These represent the factors of the ambient space $(\mathbb{P}^1)^n$. The intersection point of two lines determines a marked point on each of the two lines. For each line labeled with more than one factor, we have a compatible set of isomorphisms between those factors. Given these data, we can compute the ideal I_j of a component Y_j as follows:

1. Let X_j be the submatrix of X given by the columns indexed by F_j , acted on by the 2×2 -matrices corresponding to the isomorphisms of $\mathbb{P}^1$.
2. For each $i \notin F_j$ locate the intersection point on Y_i that is nearest to Y_j . Let $\alpha x_i + \beta y_i$ be the linear form defining this intersection point on Y_i .
3. The ideal I_j is generated by these linear forms and the 2×2 minors of X_j . The intersection ideal $I = \cap_j I_j$ is the desired point in $H_{2,n}$.

Example 4.5. The above algorithm implies that there are infinitely many $\text{PGL}(2)^n$ -orbits on $H_{2,n}$ when $n \geq 5$. Consider a tree of four lines, Y_1 , Y_2 , Y_3 , and Y_4 , which meet a fifth line in four distinct points, with coordinates $(0:1)$, $(1:1)$, $(1:0)$ and $(t:1)$ on $Y_5 = \mathbb{P}^1$. Each of these intersection points is identified with the point $V(x_j) = \{(0:1)\}$ on the line Y_j . Then we have

$$\begin{aligned} I_1 &= \langle x_2, x_3, x_4, x_5 \rangle & I_2 &= \langle x_1, x_3, x_4, x_5 - y_5 \rangle & I_3 &= \langle x_1, x_2, x_4, y_5 \rangle \\ I_4 &= \langle x_1, x_2, x_3, x_5 - ty_5 \rangle & I_5 &= \langle x_1, x_2, x_3, x_4 \rangle & I &= I_1 \cap I_2 \cap I_3 \cap I_4 \cap I_5 \end{aligned}$$

As t varies over the field K , the ideals I lie in different $\text{PGL}(2)^5$ -orbits on $H_{2,5}$ because the cross ratio of the four points on Y_5 is invariant under $\text{PGL}(2)$. $\square$

Proof of Theorem 4.1. We shall prove that $H_{2,n}$ is irreducible. Let I be any ideal in $H_{2,n}$. We use induction on the number of components of $V(I)$ to show that I is in the closure of the orbit of $I_2(X)$. If $V(I)$ is irreducible, then I is in the orbit of $I_2(X)$ by the above discussion, so we assume that $V(I)$ has at least two components. We shall construct another ideal $J \in H_{2,n}$ such that $V(J)$ has one fewer component than $V(I)$ and such that J degenerates to I .

Consider the tree picture of $V(I)$ as described above, and let Y_1 be a component which has exactly one point of intersection with the other components. Let Y_2 be one of these components intersecting Y_1 . After relabelling the factors and a change of coordinates, we can assume that the isomorphisms of the factors associated to Y_1 are all the identity map, and the same holds for the isomorphisms of factors of Y_2 . Furthermore, we can assume that the point $Y_1 \cap Y_2$ is defined in $(\mathbb{P}^1)^n$ by the ideal $\langle x_i : i \in F_1 \rangle + \langle y_i : i \notin F_1 \rangle$.

We now replace Y_2 with the component Y'_2 labeled by the set $F'_2 = F_1 \cup F_2$ (with the identity isomorphisms), and we call its ideal I' . By Proposition 4.4, I' is in $H_{2,n}$. The ideal I'_2 of Y'_2 is generated by $\{x_j : j \notin F'_2\}$ and the 2×2 minors of the submatrix of X indexed by F'_2 . For $t \neq 0$, we consider the ideal formed by replacing y_j by ty_j in I' for $j \in F_1$ and take the flat limit as t goes to 0. The limit of I'_2 under this action is $I_1 \cap I_2$, so the limit of I' is contained in I . Since I and the limit of I' have the same Hilbert function, they must be equal. This proves the first assertion in Theorem 4.1. The second assertion will follow from Corollary 4.8 below. $\square$

We now come to the combinatorial description of monomial ideals I on $H_{2,n}$. Here the tree picture can be simplified. There are precisely n components $V(I) = Y_1 \cup Y_2 \cup \dots \cup Y_n$, and the partition is into singletons $F_i = \{i\}$. Each line Y_i has only two points where intersections are possible, namely, $V(x_i) = \{(0:1)\}$ and $V(y_i) = \{(1:0)\}$. We draw Y_i as an oriented line segment with the intersection points only at the end points. The orientation is indicated by an arrow whose tail represents $V(x_i)$ and whose head represents $V(y_i)$. In this manner, each monomial ideal I in $H_{2,n}$ is represented uniquely by a tree T with n directed labeled edges. The tree T has $n+1$ vertices which remain unlabeled. This establishes the first part of Theorem 4.2.

Our construction is illustrated for $n = 3$ in Figure 1. See Example 4.10 below for a combinatorial discussion of the two classes of trees shown here.

We next describe a rule for reading off the generators of a monomial ideal $I \in H_{2,n}$ from its tree T . For any two distinct indices i and j in $\{1, \dots, n\}$ we set $z_{ij} = x_j$ if the directed edge j is pointing away from the edge i in Y




Figure 1: Edge-labelled trees corresponding to 2 of the 32 monomial ideals on $H_{2,3}$. The tree on the left corresponds to the ideal $\langle y_1y_2, y_1x_3, x_2x_3 \rangle$ and on the right corresponds to the Borel-fixed ideal $Z = \langle x_1x_2, x_1x_3, x_2x_3 \rangle$.

and $z_{ij} = y_j$ otherwise. This means that the ideal I_i of the component Y_i is generated by the variables z_{ij} for $j \in \{1, \dots, n\} \setminus \{i\}$. By intersecting these ideals for all i we obtain the following combinatorial formula for the ideal I .

Remark 4.6. The monomial ideal associated with the tree T equals

$$I = \langle z_{ij}z_{ji} : 1 \leq i < j \leq n \rangle.$$

Explicitly, the ideal generator corresponding to pair $\{i, j\}$ of edges equals

$$z_{ij}z_{ji} = \begin{cases} y_iy_j & \text{if the edges } i \text{ and } j \text{ point towards each other,} \\ x_ix_j & \text{if the edges } i \text{ and } j \text{ point away from each other,} \\ y_ix_j & \text{if the edge } i \text{ points to the edge } j \text{ but not conversely,} \\ x_iy_j & \text{if the edge } j \text{ points to the edge } i \text{ but not conversely.} \end{cases}$$

Note that the Borel-fixed ideal Z corresponds to the star tree with all edges directed outwards. Our next result concerns tangent spaces of $H_{2,n}$.

Proposition 4.7. *Let I be the monomial ideal corresponding to a directed tree T as above. Then the dimension of the tangent space of $H_{2,n}$ at I is*

$$\sum_v f(\text{degree}(v)) \tag{8}$$

where the sum is over all vertices of T , and the function f is defined by

$$f(a) = \begin{cases} 3(a-1) & \text{if } 1 \leq a \leq 3, \\ a(a-1) & \text{if } a \geq 3. \end{cases}$$

The following corollary to this result completes the proof of Theorem 4.1.

Corollary 4.8. *The monomial ideal I is a smooth point on the Hilbert scheme $H_{2,n}$ if and only if every vertex in the tree T is at most trivalent.*

Proof. The tree T has $n + 1$ vertices v , and the number of edges is

$$n = \frac{1}{2} \cdot \sum_v \text{degree}(v)$$

Since $H_{2,n}$ is a compactification of $\text{PGL}(2)^{n-1}$, its dimension equals

$$\dim(H_{2,n}) = 3(n-1) = 6n - 3(n+1) = \sum_v 3(\text{degree}(v) - 1).$$

Since $f(a) \geq 3(a-1)$, with equality if and only if $a \leq 3$, the sum in (8) is equal to $\dim(H_{2,n})$, if and only if $\text{degree}(v) \leq 3$ for all vertices v . $\square$

Example 4.9. The star tree ideal $Z = \langle x_i x_j : 1 \leq i < j \leq n \rangle$ has tangent space dimension $n(n-1)$. In fact, it is the most singular point on $H_{2,n}$, since every other ideal degenerates to Z . On the other hand, the chain ideal $M = \langle x_i y_j : 1 \leq i < j \leq n \rangle$ is a smooth point on $H_{2,n}$, because the tree for M is a chain of n directed edges. This confirms Lemma 3.4 for $d = 2$. $\square$

Proof of Proposition 4.7. For any distinct edges k and ℓ meeting at a vertex v of the tree T , we define the following tangent directions for $H_{2,n}$ at I :

$$\alpha_{k\ell}: z_{ij} z_{ji} \mapsto \begin{cases} z_{ij} \tilde{z}_{ji} & \text{if } i = k \text{ and } j \text{ connects to } v \text{ via } \ell \text{ (including } j = \ell), \\ 0 & \text{otherwise.} \end{cases}$$

Here we use the convention that $\tilde{z}_{ij} = x_j$ if $z_{ij} = y_j$ and $\tilde{z}_{ij} = y_j$ if $z_{ij} = x_j$. Moreover, if v is bivalent, i.e. k and ℓ are the only edges incident to v , define:

$$\beta_v: z_{ij} z_{ji} \mapsto \begin{cases} \tilde{z}_{ij} \tilde{z}_{ji} & \text{if } \{i, j\} = \{k, \ell\}, \\ 0 & \text{otherwise.} \end{cases}$$

If the vertex v has degree a , then we have defined $f(a)$ maps. To show that these maps are indeed tangent directions, we exhibit a one-parameter deformation of I . The tangent vector β_v is realized by the curve on $H_{2,n}$ gotten by replacing $z_{k\ell} z_{\ell k}$ with $z_{k\ell} z_{\ell k} - \epsilon \tilde{z}_{k\ell} \tilde{z}_{\ell k}$ among the generators of I . The resulting ideal in $H_{2,n}$ represents the tree of lines gotten by merging the edges k and ℓ to a single $\mathbb{P}^1$ labeled by $\{k, \ell\}$. The tangent vector $\alpha_{k\ell}$ is

realized by replacing z_{ji} with $z_{ji} - \epsilon \tilde{z}_{ji}$ in all prime components of I_j such that j connects to v via ℓ . The resulting ideal in $H_{2,n}$ represents the tree of lines gotten by sliding the subtree at v in direction ℓ along the edge labeled k .

As v ranges over all vertices of the tree v , and k, ℓ range over all incident edges, we now have a collection of tangent vectors whose cardinality equals (8). To see that these vectors are linearly independent, we note that $\alpha_{k\ell}$ is the only one of these tangent vectors such that the image of $z_{k\ell}z_{\ell k}$ has a non-zero coefficient for $z_{ij}\tilde{z}_{ji}$ and that β_v is the only one such that the image of $z_{k\ell}z_{\ell k}$ has a non-zero coefficient for $\tilde{z}_{ij}\tilde{z}_{ji}$. Thus, a non-trivial linear combination of the $\alpha_{k\ell}$ and β_v can't be the zero tangent vector.

It remains to be seen that our tangent vectors span the tangent space. Suppose there exists a tangent vector ϕ that is not in the span of the $\alpha_{k\ell}$ and β_v . After subtracting suitable multiples of these known tangent vectors, we may assume that for any pair of adjacent edges i and j there exists a scalar ν_{ij} such that $\phi(z_{ij}z_{ji}) = \nu_{ij}\tilde{z}_{ij}\tilde{z}_{ji}$, and furthermore $\nu_{ij} = 0$ if the node v shared by i and j is bivalent. Suppose that v has degree at least 3 and let k be an edge incident to v distinct from i and j . Then $z_{ij} = z_{kj}$ and hence

$$\nu_{ij}z_{jk}\tilde{z}_{ij}\tilde{z}_{ji} = z_{jk}\phi(z_{ij}z_{ji}) = \phi(z_{jk}z_{kj})z_{ji} = \nu_{jk}\tilde{z}_{jk}\tilde{z}_{kj}z_{ji}.$$

This implies $\nu_{ij} = \nu_{ik} = 0$. We conclude that $\phi(z_{ij}z_{ji}) = 0$ for any pair of adjacent edges i and j . Now suppose that i and j are not adjacent and write

$$\phi(z_{ij}z_{ji}) = \lambda z_{ij}\tilde{z}_{ji} + \mu \tilde{z}_{ij}z_{ji} + \nu \tilde{z}_{ij}\tilde{z}_{ji}.$$

Let ℓ be the edge adjacent to j on the path from i to j . Then $z_{ij} = z_{\ell j}$ and

$$0 = z_{ji}\phi(z_{\ell j}z_{j\ell}) = \phi(z_{ij}z_{ji})z_{j\ell} = 0 + \mu \tilde{z}_{ij}z_{ji}z_{j\ell} + \nu \tilde{z}_{ij}\tilde{z}_{ji}z_{j\ell} \quad \text{modulo } I.$$

This implies $\mu = \nu = 0$ and, by symmetry, $\lambda = 0$. We conclude that $\phi = 0$, so our maps $\alpha_{k\ell}$ and β_v form a basis for the tangent space of $H_{2,n}$ at I . $\square$

Proof of Theorem 4.2. We already saw that the monomial ideals on $H_{2,n}$ are in bijection with trees T with $n+1$ unlabeled vertices and directed edges that are labeled with $\{1, \dots, n\}$. To show that there are $2^n(n+1)^{n-2}$ monomial ideals, it suffices to show there are $(n+1)^{n-2}$ edge-labeled trees on $n+1$ vertices. Picking an arbitrary node as the root and shifting the labels to the nodes away from the root gives a rooted, node-labeled tree, of which there are $(n+1)^{n-1}$. From the rooted tree, we can uniquely recover the edge-labeled tree and the choice of the root, so there are $(n+1)^{n-2}$ edge-labeled trees.

We now need to identify all ideals I in $H_{2,n}$ that possess precisely two initial monomial ideals. We already saw two classes of such ideals in the proof of Proposition 4.7. First, there was the ideal with generator $z_{k\ell}z_{\ell k} - \epsilon\tilde{z}_{k\ell}\tilde{z}_{\ell k}$ which realizes the deformation β_v and swap # 2 in the statement of Theorem 4.2. We also exhibited an ideal for the deformation $\alpha_{k\ell}$ which realizes the move # 1 when the subset of trees is a singleton. The general case is subsumed by the following argument.

In light of Theorem 2.1 and Proposition 3.1, Gröbner degenerations of ideals I in $H_{2,n}$ correspond to scheme-theoretic limits of the trees Y with respect to one-parameter subgroups of the $(K^*)^n$ -action on $(\mathbb{P}^1)^n$.

Let Y be a tree on $H_{2,n}$ that has precisely two degenerations to $(K^*)^n$ -fixed trees. There are two cases to be considered. First suppose that some component Y_i of Y is labeled by a subset $F_i \subseteq \{1, \dots, n\}$ with $|F_i| \geq 2$. The component Y_i admits $|F_i|!$ distinct degenerations to a $(K^*)^n$ -fixed tree, and, by Proposition 4.4, each of these lifts to a degeneration of Y . This implies that the tree Y has $n - 1$ edges, and the unique non-singleton label F_i has cardinality two. Moreover, each intersection point $Y_j \cap Y_k$ is a torus-fixed point on both Y_j and Y_k . This is precisely the situation in swap # 2 above.

In the second case to be considered, the tree Y consists of the n lines $Y_1, \dots, Y_n$, each labeled by a singleton. Consider all components Y_i with intersection points that are not torus-fixed. For each such component Y_i there exist two torus degenerations of Y that move the intersection points on Y_i to the two torus-fixed points on Y_i . Under these degenerations, the intersection points $Y_j \cap Y_k$ with $i \notin \{j, k\}$ remain in their positions on Y_j and Y_k . Hence, only one component Y_i has intersection points that are not torus-fixed. This is precisely the situation in move # 1 in Theorem 4.2. $\square$

Example 4.10. ($n = 3$) The Hilbert scheme $H_{2,3}$ has 32 monomial ideals, corresponding to the eight orientations on the claw tree and to the eight orientations on each of the three labeled bivalent trees. Representatives for the two classes of trees are shown in Figure 1. The eight orientations of the claw tree can be arranged into the vertices of a cube. Each edge in the cube is an edge in the graph corresponding to moving two edges at a time between vertices. Along each edge, add two vertices corresponding to bivalent trees and four edges from each of these to each adjacent vertex of the cube. In addition to these operations corresponding to move # 1, there are 24 edges corresponding to swap # 2. These are arranged into four hexagons.

The six-dimensional manifold $H_{2,3}$ coincides with Lafforgue's compactification $X_{2,3}$ in Remark 3.8. Here, (6) amounts to an embedding of $H_{2,3}$ into $\mathbb{P}^3 \times \mathbb{P}^3 \times \mathbb{P}^3$. The equations for this embedding are as follows. Each $\mathbb{P}^3$ parametrizes one of the three generators $a_{ij}x_ix_j + b_{ij}x_iy_j + c_{ij}y_ix_j + d_{ij}y_iy_j$, $1 \leq i < j \leq 3$, of an ideal in $H_{2,3}$. Being a point in $H_{2,3}$ means that these ideal generators admit two linearly independent syzygies in degree $(1, 1, 1)$. This happens if and only if the following 6×9 -matrix has rank at most four:

$$\begin{pmatrix} a_{12} & 0 & b_{12} & 0 & c_{12} & 0 & d_{12} & 0 \\ 0 & a_{12} & 0 & b_{12} & 0 & c_{12} & 0 & d_{12} \\ a_{13} & b_{13} & 0 & 0 & c_{13} & d_{13} & 0 & 0 \\ 0 & 0 & a_{13} & b_{13} & 0 & 0 & c_{13} & d_{13} \\ a_{23} & b_{23} & c_{23} & d_{23} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & a_{23} & b_{23} & c_{23} & d_{23} \end{pmatrix}$$

By saturating its ideal of 5×5 -minors with respect to the irrelevant ideal $\bigcap_{1 \leq i < j \leq 3} \langle a_{ij}, b_{ij}, c_{ij}, d_{ij} \rangle$, we find that the prime ideal of $X_{2,3} \subset (\mathbb{P}^3)^3$ is generated by nine cubics such as $a_{12}a_{13}d_{23} - a_{12}b_{13}c_{23} - b_{12}a_{13}b_{23} + b_{12}b_{13}a_{23}$. $\square$

5 Three projective planes

In this section we study the smallest case where $H_{d,n}$ is reducible, namely, $n = d = 3$. The multigraded Hilbert scheme $H_{3,3}$ parametrizes degenerations of the projective plane in its diagonal embedding $\mathbb{P}^2 \hookrightarrow \mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^2$. We use the following notation for the unknowns x_{ij} in the polynomial ring $K[X]$.

$$X = \begin{bmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{bmatrix}$$

Theorem 5.1. *The multigraded Hilbert scheme $H_{3,3}$ is the reduced union of seven irreducible components, each of which contains a dense $\mathrm{PGL}(3)^3$ orbit:*

- The 16-dimensional main component $\overline{\mathrm{PGL}(3)^3/\mathrm{PGL}(3)}$ is singular.
- Three 14-dimensional smooth components are permuted under the S_3 -action on $(\mathbb{P}^2)^3$. At a generic point, the subscheme of $(\mathbb{P}^2)^3$ is the union of the blow-up of $\mathbb{P}^2$ at a point, two copies of $\mathbb{P}^2$, and $\mathbb{P}^1 \times \mathbb{P}^1$. An ideal

which represents such a point on this component is

$$\begin{aligned} & \langle x_1, x_2, y_1 z_2 - z_1 y_2, y_1 y_3 - z_1 x_3, y_2 y_3 - z_2 x_3 \rangle \cap \\ & \langle x_1, y_1, x_3, y_3 \rangle \cap \langle x_1, x_2, x_3, y_3 \rangle \cap \langle x_2, y_2, x_3, y_3 \rangle. \end{aligned} \quad (9)$$

- Three 13-dimensional smooth components are permuted under the S_3 -action on $(\mathbb{P}^2)^3$. A generic point looks like the union of three copies of $\mathbb{P}^2$ and $\mathbb{P}^2$ blown up at three points. A representative ideal is

$$\langle x_1, y_1, x_2, z_2 \rangle \cap \langle x_1, y_1, x_3, y_3 \rangle \cap \langle x_2, y_2, x_3, y_3 \rangle \cap \langle x_1, x_2, x_3, y_1 y_2 z_3 - z_1 z_2 y_3 \rangle.$$

With some additional notation, we can describe the isomorphism types of the six extra components. Let Fl denote the variety of complete flags in K^3 and $\mathcal{O}_i$ the tautological bundle of i -dimensional vector spaces for $i = 1$ or 2 . Then the second class of components are isomorphic to the bundle

$$\mathcal{H}_{2,3}(\mathbb{P}^2 \rightarrow \mathbb{P}(\mathcal{O}_2) \times \mathbb{P}(\mathcal{O}_2) \times \mathbb{P}(\mathcal{O}^{\oplus 3}/\mathcal{O}(-1))) \rightarrow \text{Fl} \times \text{Fl} \times \mathbb{P}^2 \quad (10)$$

where $\mathcal{H}_{2,3}$ is a bundle whose fibers are each isomorphic to the Hilbert scheme $H_{2,3}$. A point in this bundle is equivalent to a point x in $\text{Fl} \times \text{Fl} \times \mathbb{P}^2$, together with an ideal in the total coordinate ring of $\mathbb{P}((\mathcal{O}_2)_x) \times \mathbb{P}((\mathcal{O}_2)_x) \times \mathbb{P}((\mathcal{O}^{\oplus 3}/\mathcal{O}(-1))_x)$ with the appropriate Hilbert function (1).

To relate this formulation to the ideal in (9), we identify linear forms in $K[X]$ with the direct sum of three vector spaces, each of dimension 3, and $\{x_i, y_i, z_i\}$ as choice of basis for the i th summand. The two flag varieties parametrize the duals of the flags $\langle x_i \rangle \subset \langle x_i, y_i \rangle$ for $i = 1, 2$. The projective space parametrizes the point whose ideal is $\langle x_3, y_3 \rangle$. These spaces determine all the linear generators in (9). The additional generators of the first component represent a point in $H_{2,3}$, but without a canonical choice of basis.

The third class of components are isomorphic to the projective bundle:

$$\mathbb{P}(\mathcal{E}) \rightarrow \text{Fl} \times B \times \text{Fl}$$

where Fl is as before and B is the blow-up of $\mathbb{P}^2 \times \mathbb{P}^2$ along the diagonal. We think of the blow-up variety B as the parameter space of two points in $\mathbb{P}^2$ and a line containing them. The 4-dimensional vector bundle $\mathcal{E}$ is the sum

$$(\mathcal{O}_1 \otimes \mathcal{O}_1 \otimes \mathcal{O}_2) + (\mathcal{O}_1 \otimes \mathcal{O}_2 \otimes \mathcal{O}_1) + (\mathcal{O}_2 \otimes \mathcal{O}'_1 \otimes \mathcal{O}_1) \quad (11)$$

inside $\mathcal{O}_{\mathbb{P}^1}^{\oplus 3} \otimes \mathcal{O}_B^{\oplus 3} \otimes \mathcal{O}_{\mathbb{P}^1}^{\oplus 3}$, where $\mathcal{O}_1$ and $\mathcal{O}'_1$ are the pullbacks to B of $\mathcal{O}(-1)$ on each of the copies of $\mathbb{P}^2$, which parametrize the two points. The flag varieties parametrize the duals of $\langle x_i \rangle \subset \langle x_i, y_i \rangle$ for $i = 1, 3$ and B parametrizes the two points defined by $\langle x_2, y_2 \rangle$ and $\langle x_2, z_2 \rangle$, with $\langle x_2 \rangle$ as the line between them. As before, these vector spaces determine the linear generators of the components. The bundle $\mathbb{P}(\mathcal{E})$ parametrizes the coefficients of the cubic generator of the ideal of the blowup of $\mathbb{P}^2$ at two points. This ideal equals

$$\langle x_1, x_2, x_3, ay_1y_2z_3 + by_1y_2y_3 + cy_1z_2y_3 + dz_1z_2y_3 \rangle \quad (12)$$

Note that the middle two terms are linearly independent even when y_2 and z_2 coincide. For generic coordinates, after a change of basis, we can take b and c to be zero, and after rescaling, we take $a = d = 1$. Thus, the $\mathrm{PGL}(3)^3$ orbit of the ideal (5.1) is dense in the 13-dimensional component of $H_{3,3}$.

Proof of Theorem 5.1: The proof is computational. It rests on the **Singular** code posted at <http://math.berkeley.edu/~dustin/diagonal/h33.sng>. The computation works regardless of what the field characteristic of K is.

It suffices to consider an affine neighborhood of the unique Borel-fixed ideal Z in $H_{3,3}$. We employ the standard method of choosing coordinates by adding trailing terms with indeterminate coefficients to the ten monomial generators of Z . The ideal defining $H_{3,3}$ is then derived from the syzygies of Z . We then derive the prime ideals representing each of the seven components, by translating the geometric descriptions above into local coordinates around Z . Implicitization using **Singular** yields the seven prime ideals, and we check that their intersection equals the ideal of the Hilbert scheme itself.

We now explain how the parametric representations of the seven components are derived. The main component is, by definition, parametrized by the $\mathrm{PGL}(3)^3$ orbit of the ideal of 2×2 minors of X . The other components can also be parametrized by $\mathrm{PGL}(3)^3$ orbits of the representative ideals, but it is also possible – and computationally more efficient – to use parametrizations which do not require localization.

For the 14-dimensional components, we begin by using local coordinates in $H_{2,3}$ to define a family of subschemes of $(\mathbb{P}^1)^3$ over $\mathbb{A}^6$. Renaming some of the variables and adding two linear terms, we get the parametrization of the blow-up of $\mathbb{P}^2$ and its degenerations in $(\mathbb{P}^2)^3$, i.e. a neighborhood of a fiber of (10). Intersecting with the ideals of linear spaces gives the family in $K[X]$ with the appropriate Hilbert function. Different choices of flags can be represented by upper triangular changes of coordinates on each of the

	T. sp.	planar?	main?	component		symm.
				14-dim.?	13-dim.?	
1	16	y	y	n	n	2
2	16	y	y	n	n	1
3	16	y	y	n	n	1
4	18	y	y	n	n	6
5	16	y	y	n	n	3
6	14	y	n	y	n	2
7	15	y	n	y	y	1
8	16	n	y	n	n	1
9	17	n	y	y	n	1
10	18	n	y	n	n	2
11	17	n	y	y	n	1
12	14	n	n	y	n	2
13	18	n	y	y	y	2
14	18	n	y	y	n	2
15	18	n	y	y	y	1
16	18	n	y	y	y	6

Table 1: The symmetry classes of monomial ideals in $H_{3,3}$

three columns of X . The parametrization of the 13-dimensional component follows the same pattern. In a neighborhood of Z , up to change of basis, we can take the two points parametrized by B to be $\langle x_2, y_2 \rangle$ and $\langle x_2, y_2 - az_3 \rangle$ with $\langle x_2 \rangle$ as the line between them. In addition to a , the other coordinates are the entries of the upper triangular matrix corresponding to the choice of flag and to the coefficients of the cubic generator in (12). Implicitizing these parametrizations reveals the prime ideals for these seven components, and their intersection is found to equal the ideal of the Hilbert scheme itself. $\square$

In Table 1 we show that $H_{3,3}$ contains 13824 monomial ideals. They come in 16 symmetry classes, and we list them in four groups, corresponding to the dimension (12, 11, 10, and 9) of the orbit under the action of $\text{PGL}(3)^3$. The 16 monomial ideals appear in the same order as their pictorial representation in Figure 2. The third column indicates whether or not the picture is planar. The second column is the tangent space dimension of $H_{3,3}$ at that point. The triple column shows which components the monomial ideals live on. The rightmost column shows the order of the symmetry group of the ideal. Note

that the permutation group acting on $H_{3,3}$ has order $6^4 = 1296$: it permutes the three factors of $\mathbb{P}^2 \times \mathbb{P}^2 \times \mathbb{P}^2$ as well as the three coordinates $\{x_i, y_i, z_i\}$ of each projective plane. The total number 13824 of monomial ideals on $H_{3,3}$ equals 1296 times the sum of the reciprocals in the last column of Table 1.

Every monomial ideal in $H_{3,3}$ corresponds to a polyhedral complex in the boundary in the direct product of three triangles, denoted $(\Delta_2)^3$. Using the moment map of toric geometry, each such polyhedral complex can be identified with the real positive points of the corresponding subscheme of $(\mathbb{P}^2)^3$. The following conditions characterize those subcomplexes of $H_{3,3}$ whose corresponding monomial ideal has the right multigraded Hilbert function:

- For every vector of non-negative integers (t_1, t_2, t_3) summing to 2, there is exactly one 2-dimensional cell consisting of the product of a t_1 -dimensional cell of Δ_2 , a t_2 -dimensional cell, and a t_3 -dimensional cell.
- The complex contains exactly ten 0-cells, 15 1-cells, and six 2-cells.

This characterization is sufficient to show that Table 1 is complete. By the condition on the number of 1-cells, each triangle must meet at least two of the squares. Thus, each pair of squares must be adjacent or be connected by a triangle, and so all three squares must meet in a common point. There are four possible configurations of the squares: either 0, 1, 2, or 3 edges common to multiple squares. The monomial ideals can be enumerated by considering all possible ways to attach the additional triangles to these configurations.

The monomial ideals form a poset based on containment within the closures of their orbits, illustrated in Figure 2. Each ideal is drawn as a 2-dimensional subcomplex of $(\Delta^2)^3$. The subcomplex is drawn abstractly, but the embedding amounts to a choice of labellings. The bold lines indicate that an additional triangle is attached along that edge. By orbits, we mean orbits under the disconnected group which is generated by multiplying the first column by an arbitrary matrix and by the discrete action of permuting the columns. The number on the lower right is the dimension of the tangent space of $H_{3,3}$ at that monomial ideal. The ranking is by the dimension of (every component of) the orbit of the monomial ideal: 9, 10, 11, or 12.

The maximal elements of the poset in Figure 2 correspond to the “planar” complexes, i.e. those such that no edge contains more than two 2-cells. By fixing an isomorphism between the three copies of Δ_2 , we get a projection from $(\Delta_2)^3$ onto $3\Delta_2$. In the case of ideals 1, 2, 3, 4, and 5 the corresponding




Figure 2: Partial ordering of the monomial ideals on $H_{3,3}$

subcomplexes project to tilings of $3\Delta_2$. In these cases, there is a monomial ideal in the orbit which is in the toric Hilbert scheme of $\Delta_2 \times \Delta_2$, and the corresponding triangulation of $\Delta_2 \times \Delta_2$ is related to the tiling of $3\Delta_2$ by the Cayley trick. The tilings in [15, Figure 5] correspond to the monomial ideals 1, 3, 4, 5, and 2 in this order. Each triangulation is regular, and the ideals are smoothable, even in the toric Hilbert scheme. Here, the toric Hilbert is the subscheme of $H_{3,3}$ obtained by fixing the Hilbert function of $I_2(X)$ with respect to the finer grading given by both row degrees and column degrees. For details, references and further information see [8, §2] and [15, Theorem 2].

6 Deligne schemes and their special fibers

The original motivation which started this project was a discussion with Annette Werner about tropical convexity, and its connection to affine buildings and moduli of hyperplane arrangements as developed by Keel and Tevelev [11]. Our aim was to understand the Deligne schemes of [11, §1] and their special fibers in the concrete language of combinatorial commutative algebra. We found that the multigraded Hilbert scheme $H_{d,n}$ offers a suitable frame-

work for studying Deligne schemes and their arithmetic. In this section we briefly discuss the set-up and the connection to the combinatorial results in [2, 9]. We plan to pursue this further in a joint project with Annette Werner.

Let K be an algebraically closed field with a non-trivial non-archimedean absolute value, let k be the residue field of K , and R the valuation ring of K . In computational studies (such as [9]) we usually relax the requirement that K be algebraically closed, and we work with the Gröbner-friendly scenario $K = \mathbb{Q}(z)$, $R = \mathbb{Q}[z]$ and $k = \mathbb{Q}$. Let $\mathcal{B}$ denote the Bruhat-Tits building associated with the group $\mathrm{PGL}(d)$ over K as defined in [9, 11]. The building $\mathcal{B}$ is an infinite simplicial complex of dimension $d - 1$ whose vertices are the equivalence classes of R -submodules of K^d having maximal rank d .

Let $Y = \{Y_1, \dots, Y_n\}$ be a finite set of vertices of the affine building $\mathcal{B}$. Following [11, Definition 1.8], we let $\mathbb{S}_Y$ denote the corresponding join of projective spaces over R , and we write S_Y for its special fiber over k . In the special case when Y is a convex subset of $\mathcal{B}$, a classical result of Mustafin [14] states that $\mathbb{S}_Y$ is semi-stable over R , which implies that S_Y has smooth irreducible components with normal crossings. In this section we allow Y to be any finite set of vertices – not necessarily convex – of the building $\mathcal{B}$. Following [11, 1.10], we shall call $\mathbb{S}_Y$ the *Deligne scheme* of the subset $Y \subset \mathcal{B}$.

We now describe the Deligne scheme $\mathbb{S}_Y$ and its special fiber S_Y in concrete terms. The configuration Y is represented by $(Y_1, Y_2, \dots, Y_n)$, an n -tuple of invertible $d \times d$ -matrices with entries in the field K . This data is the input for the algorithm of [9] which computes the convex hull of Y in $\mathcal{B}$. Let $I_2(X)$ be the ideal of 2×2 -minors of a $d \times n$ -matrix of unknowns. We consider the transformed ideal $Y \circ I_2(X)$ in $K[X]$, and we intersect it with $R[X]$:

$$\mathbb{I}_Y = (Y \circ I_2(X)) \cap R[X]. \quad (13)$$

We call $\mathbb{I}_Y \subset R[X]$ the *Deligne ideal* of the point configuration $Y \subset \mathcal{B}$. The image of $\mathbb{I}_Y$ under the specialization $R \rightarrow k$ is denoted $I_Y \subset k[X]$. We call I_Y the *special fiber ideal* of Y . This nomenclature is justified as follows.

Remark 6.1. The Deligne scheme $\mathbb{S}_Y$ coincides with the subscheme of $(\mathbb{P}_R^{d-1})^n$ defined by the Deligne ideal $\mathbb{I}_Y$, and its special fiber S_Y coincides with the subscheme of $(\mathbb{P}_k^{d-1})^n$ defined by the special fiber ideal I_Y .

Remark 6.1 implies that the Deligne scheme $\mathbb{S}_Y$ is a point in the multi-graded Hilbert scheme $H_{d,n}(R)$ over the valuation ring R , and its special fiber S_Y is a point in $H_{d,n}(k)$ over its residue field k . Thus our study in Sections 2 to 5 is relevant for Deligne schemes. In particular, Theorem 2.1 implies:

Corollary 6.2. *The special fiber S_Y of the Deligne scheme $\mathbb{S}_Y$ is reduced.*

The formula (13) translates into the following Gröbner-based algorithm for computing Deligne schemes and their special fibers when $K = \mathbb{Q}(z)$, $R = \mathbb{Q}[z]$, $k = \mathbb{Q}$. The input data is an n -tuple $Y(z)$ of invertible $d \times d$ -matrices whose entries are rational functions in one variable z . The Deligne ideal $\mathbb{I}_{Y(z)}$ is an ideal in the polynomial ring $\mathbb{Q}[z, X]$. It is computed as follows. We replace the j -th column of the matrix $X = (x_{ij})$ by its left multiplication with the j -th input matrix $Y_j(z)$. We then form the 2×2 -minors of the resulting matrix and multiply each generator by a power of z to get an ideal L in $\mathbb{Q}[z, X]$. The Deligne ideal is then obtained by saturation as follows:

$$\mathbb{I}_{Y(z)} = (L : \langle z \rangle^\infty). \quad (14)$$

The special fiber ideal is obtained by setting z to zero in the Deligne ideal:

$$I_{Y(z)} = \mathbb{I}_{Y(z)}|_{z=0}. \quad (15)$$

This algorithm generalizes the construction of Block and Yu [2] which concerns the special case when the configuration $Y(z)$ lies in one apartment of the Bruhat-Tits building $\mathcal{B}$. Algebraically, this means that the $Y_j(z)$ are diagonal matrices, and, geometrically, the apartment of $\mathcal{B}$ is identified with the standard triangulation of tropical projective space $\mathbb{TP}^{d-1}$. If the diagonal entries of the $Y_j(z)$ are monomials with sufficiently generic exponents, then (14)-(15) simply amounts to a Gröbner basis computation for the ideal $I_2(X)$. Block and Yu [2] consider the Alexander dual of the initial monomial ideal of $I_2(X)$, and they showed that minimal free resolution of that Alexander dual is cellular. It represents the convex hull of $Y(z)$ in $\mathbb{TP}^{d-1}$, and hence in $\mathcal{B}$. We conjecture that this method can be adapted to compute the convex hull of $Y(z)$ even if the matrices $Y_j(z)$ do not lie in a common apartment of $\mathcal{B}$.

We note that the algorithm (14)-(15) is not very practical for computing the special fiber ideal $I_{Y(z)}$. We found that running the saturation step (14) in naive manner in `Macaulay 2` is too slow for interesting values of d and n .

Instead, we propose the following approach. If the $Y(z)$ are “sufficiently generic,” then we replace each $Y_j(z)$ by a “nearby” matrix of the special form

$$Y_j(z) \approx \text{diag}(z^{w_{1j}}, \dots, z^{w_{dj}}) \cdot A_i. \quad (16)$$

Here the weights w_{ij} are generic rational numbers that specify a term order $>$ on the polynomial ring $\mathbb{Q}[X]$, and $A = (A_1, \dots, A_n)$ is a tuple of invertible

matrices over $\mathbb{Q}$. The precise meaning of the approximation (16) is that

$$I_{Y(z)} = \text{in}_>(A \circ I_2(X)). \quad (17)$$

When this holds the special fiber $S_{Y(z)}$ of the Deligne scheme $\mathbb{S}_{Y(z)}$ is given by a squarefree monomial ideal $I_{Y(z)}$. The point is that the right hand side of (17) can be computed much faster in practice than evaluating (14). It amounts to computing a Gröbner basis of the transformed ideal $A \circ I_2(X)$ in the polynomial ring $\mathbb{Q}[X]$. When the A_i are diagonal matrices over $\mathbb{Q}$ this is precisely the algorithm of Block and Yu [2] for convex hulls in $\mathbb{TP}^{d-1}$.

Acknowledgements. Both authors were supported by the U.S. National Science Foundation (DMS-0354321, DMS-0456960 and DMS-0757236). We thank Michel Brion, Aldo Conca and Annette Werner for helpful comments.

References

- [1] K. Altmann and B. Sturmfels: The graph of monomial ideals, *Journal of Pure and Applied Algebra* **201** (2005) 250–263.
- [2] F. Block and J. Yu: Tropical convexity via cellular resolutions, *Journal of Algebraic Combinatorics* **24** (2006) 103–114.
- [3] M. Brion: Group completions via Hilbert schemes, *Journal of Algebraic Geometry* **12** (2003) 605–626.
- [4] M. Brion: Multiplicity-free subvarieties, *Contemporary Math.* **331** (2003) 13–23.
- [5] A. Conca: Linear spaces, transversal polymatroids and ASL domains, *Journal of Algebraic Combinatorics* **25** (2007) 25–41.
- [6] D. Eisenbud: *Commutative Algebra With a View Toward Algebraic Geometry*, Graduate Texts in Mathematics **150**, Springer, New York, 1995.
- [7] V. Guillemin, T. Holm and C. Zara: A GKM description of the equivariant cohomology ring of a homogeneous space, *Journal of Algebraic Combinatorics* **23** (2006) 21–41.
- [8] M. Haiman and B. Sturmfels: Multigraded Hilbert schemes, *Journal of Algebraic Geometry* **13** (2004) 725–769.

- [9] M. Joswig, B. Sturmfels and J. Yu: Affine buildings and tropical convexity, *Albanian Journal of Mathematics* **1** (2007) 187–211.
- [10] A. Gibney and D. Maclagan: Equations for Chow and Hilbert quotients, [arXiv:0707.1801](#).
- [11] S. Keel and J. Tevelev: Geometry of Chow quotients of Grassmannians, *Duke Mathematical Journal* **134** (2006) 259–311.
- [12] L. Lafforgue: Pavages des simplexes, schémas de graphes recollés et compactification des $\mathrm{PGL}_r^{n+1}/\mathrm{PGL}_r$, *Invent. math.* **136** (1999) 233–271; Erratum in *Invent. math.* **145** (2001) 619–610.
- [13] E. Miller and B. Sturmfels: *Combinatorial Commutative Algebra*, Graduate Texts in Mathematics **227**, Springer, New York, 2004.
- [14] G.A. Mustafin: Non-archimedean uniformization, *Math. USSR Sbornik* **34** (1978) 187–214.
- [15] F. Santos: The Cayley trick and triangulations of products of simplices, In *Integer Points in Polyhedra - Geometry, Number Theory, Algebra, Optimization*, (eds. A. Barvinok, M. Beck, C. Haase, B. Reznick, V. Welker), Contemporary Mathematics **374**, AMS, Providence, 2005.
- [16] R. Stanley: *Combinatorics and Commutative Algebra*, Second Edition, Birkhäuser, Boston, 2004.
- [17] M. Thaddeus: Complete collineations revisited, *Mathematische Annalen* **315** (1999) 469–495.

Authors’ addresses:

Dustin Cartwright and Bernd Sturmfels, Dept. of Mathematics, University of California, Berkeley, CA 94720, USA, {dustin,bernd}@math.berkeley.edu